



In vitro* antimycotic study of essential oils against lipophilic yeast like fungus, *Malassezia

Meenakshi Sharma and Richa Sharma*

Laboratory of Mycology and Microbiology, Department of Botany, University of Rajasthan, Jaipur-302004, India

*e-mail: richa.phd.15@gmail.com

(Received: November 22, 2009; Revised received: January 28, 2010; Accepted: February 04, 2010)

Abstract: Potential inhibitory effect of *Citrus lemon* (lemon) and *Citrus sinensis* (orange) on lipophilic, yeast like fungus, *Malassezia furfur* which causes *Pityriasis versicolor*, chronic superficial fungal disease of the skin have been studied by using two different methods: disc diffusion and microdilution methods. In screening of lemon and orange oil by disc diffusion method, the diameter of inhibition zone was found to be 50 and 20 mm which was greater than inhibition zone of reference antibiotics *i.e.* gentamycin and streptomycin of 16.5 and 17 mm respectively. Minimum inhibitory concentration (MIC) of lemon and orange oil against *Malassezia furfur* was found to be 0.8 and 2.2 $\mu\text{l ml}^{-1}$. These findings support the use of *Citrus lemon* (lemon) and *Citrus sinensis* (orange) oil as a traditional herbal medicine for the control of *Pityriasis versicolor* infection of skin.

Key words: *Pityriasis versicolor*, Skin, *Malassezia furfur*, *Citrus lemon* and *Citrus sinensis* oils

Introduction

Pityriasis versicolor (PV) is a superficial mycosis, affecting the superficial layer of stratum corneum (Marcon and Powell, 1992). The causative organism is *Malassezia furfur*, a yeast like lipophilic fungus. Previously the mycelial form was called either *Pityrosporum ovale* or *Pityrosporum orbiculare* (Hay and Moore, 1998). In 1951, Gordon isolated the organism *M. furfur* and renamed it *P. orbiculare*. It was recognized that *M. furfur* is the correct name and that *P. orbiculare* and *P. ovale* are synonyms (Silva-Lizama, 1995). The disease is most prevalent in early adulthood and small childrens are rarely affected (Boussida *et al.*, 1998). PV is common in the post-pubertal age where sebaceous glands are active and in individuals who sweat more (Schmidt, 1997). There is often a positive family history of the disease. An increase in humidity, temperature and hyperhidrosis are important predisposing factors (Silva-Lizama, 1995; Boussida *et al.*, 1998). The prevalence in colder climates is less than 1% (Rippon, 1982). *M. furfur* is a component of normal skin flora in more than 90% of adults living in tropical areas (Faergemann and Freidriksson, 1981). PV, consequently, is more common in the tropics than in temperate zones (Hay and Moore, 1998).

Infectious diseases accounts for high proportion of health problems in the developing countries including India. Microorganisms have developed resistance to many antibiotics and as a result, immense clinical problem in the treatment of infectious diseases has been created (Davies, 1994). The resistance of the organisms increased due to indiscriminate use of commercial antimicrobial drugs commonly used for the treatment of infectious disease. This situation forced the researchers to search for new antimicrobial substance from various sources including medicinal plants (Bauer *et al.*, 1996). There are alarming reports of opportunistic fungal infections (Singh, 2001). A survey of literature reveals that there are many essential oils which possesses antifungal activity. Therefore, we need to search plant derived antifungal drugs which are safe and without side-effects. The herbal medicines may be in form of powders, liquids, or mixtures, which may be raw or boiled, ointments, liniments, and incisions (Alex *et al.*, 1998). Essential oils

have been screened for their potential uses as alternative remedies for the treatment of many infectious diseases (Sofowora, 1993). Essential oils have been shown to possess antibacterial, antifungal, antiviral, insecticidal and antioxidant properties (Bektas *et al.*, 2004; Burt, 2004). Numerous essential oils have been tested for both *in vitro* as well as *in vivo* antifungal activity and some pose much potential as antifungal agents. Studies on the antimicrobial, especially antibacterial and antifungal activity of lemon grass oil and its components were reported (Kordali *et al.*, 2005; Paranagama *et al.*, 2003). Orange oil was composed principally of d-limonene, 94%, and myrcene at about 3%. The essential oils of lemon and orange exhibited antifungal activity against common moulds *A. niger*, *A. flavus*, *Penicillium chrysogenum* and *P. verrucosum* (Viuda-Martos *et al.*, 2008).

In the present study, antifungal activities of lemon and orange oils were investigated with the aim to discover the medicinal potential of these oils for future application as an antifungal agent for *Pityriasis versicolor* disease.

Materials and Methods

Extraction and preparation of oil: In summer season, extraction of oil from the peels of lemon (*Citrus lemon*) and orange (*Citrus sinensis*) were carried out by standard hydrodistillation method, Clevenger's apparatus and all operations were carried out at room temperature. The fresh peels of lemon and orange were washed to remove soil and sliced. Sliced pieces of fresh peels of lemon and orange (250 g) were placed in a separate flask, together with distilled water (1L). After 5-6 hr, oil was collected from the apparatus, anhydrous with sodium sulphate for removal of water traces, then this 100% pure essential oil were dispensed into dark bottles and stored at 4°C until used. Essential oil was ready to use for disc diffusion test and determination of MIC.

Screening of essential oil using disc diffusion method: Oil was screened for their antifungal activity against *M. furfur* by disc diffusion method (Rios *et al.*, 1988). Standard size Whatman No. 1 filter paper discs, 6.0 mm in diameter, sterilized by dry heat at 140°C in an oven for one hour were used to determine antifungal

activity. SDA medium for disc diffusion test was prepared. After sterilization, it was poured into sterilized petri plates and allowed to solidify. Then one day old, fresh culture of yeast will be used for inoculum preparation. A suspension that was just turbid (~0.5 McFarland standard) by visual inspection was prepared by suspending yeast in 0.9% NaCl solution and the homogeneous suspension was used for inoculation. Using a sterile cotton swab, yeast culture were swabbed on the surface of sterile Sabourauds dextrose agar plates. Sterilized filter paper discs were soaked in neat, undiluted (100%) concentration of oils. Using an ethanol dipped and flamed forceps, oil saturated discs of 100% concentration were aseptically placed over Sabourauds dextrose agar plates seeded with the respective test microorganism. The antibiotic discs of gentamycin (30 mcg), streptomycin, clotrimazole and ketoconazole (10 mcg disc⁻¹) were also aseptically placed over the seeded Sabourauds dextrose agar plates as a standard drugs for comparison of antifungal activity of lemon and orange oils. The plates were incubated at 37°C for 24 hr. The diameter of the inhibition zones was measured in mm. Three replicates were kept in each case and average values were calculated. The activity of oils was measured by the following formula.

AI (Activity index) = Inhibition zone of sample / Inhibition zone of standard

Determination of minimum inhibitory concentration using microdilution method: The modified microdilution method Provine and Hadley (2000) was followed to determine MIC. Media used for MIC was semisolid agar media (Brain Heart Infusion Agar) aliquots of semisolid agar media (Bacto Agar; Difco Laboratories) at a pH of approximately 7.4 were poured into a 16- by 125-mm glass tubes and autoclaved. A suspension that was just turbid (~0.5 McFarland standard) by visual inspection was prepared by suspending the

selected fungi in 0.9% NaCl solution, vortexing, and homogeneous suspension was used for inoculation. Different concentrations of lemon and orange oils were added in media containing test-tubes, afterwards a standard platinum loopful (~0.001 ml, Himedia, Flexiloop) of the inoculum suspension was inserted deep into each tube of medium containing a different concentration of oils, as well as a oil-free control, by a centered down-up motion to form a two dimensional inoculum. The tubes were then incubated at 30°C for 48 hr to determine the MIC. MIC was read to be the lowest concentration at which there was no visible growth of the organism. Then, by visual inspection, good growth of the respective fungi in oil-free medium as a control was detected (48 hr for yeasts). Afterwards, the growth in all tubes at different concentrations of lemon and orange oil were compared with that of the oil-free control in order to determine inhibition.

Results

In our present work, *M. furfur* was found main etiological agent of *Pityriasis versicolor* disease hence the antimycotic studies was carried out on *M. furfur*. The conventional treatment of fungal disease is limited, and part of the reason is due to the limited spectrum of the currently antifungal drugs, and the expensive treatment, particularly due to the need of prolonged therapy. Thus, new drugs and alternative therapies are necessary, including natural products. We report here the antimycotic study of lemon and orange oil against *M. furfur in vitro*. The results of the present work on the antifungal activity of lemon and orange oil against *M. furfur* studied by two methods i.e disc diffusion and microdilution method are presented in Table 1, 2, 3 and Fig. 1. In our study lemon and orange oils presented higher diameter of inhibition zones than gentamycin, streptomycin, clotrimazole and ketoconazole. The diameter of the inhibition zone




Fig. 1: Comparative efficacy of *Citrus lemon* (lemon) and *Citrus sinensis* (orange) oil with standard allopathic drugs against *Malassezia furfur* (a) Inhibition zone at 100% concentration of lemon oil (b) Orange oil (c) Inhibition zone of streptomycin (d) Gentamycin

Table - 1: Antifungal activity of lemon and orange oil against *Malassezia furfur*

Oil	Test strain	Concentration of lemon and orange oil	IZ of sample	IZ of standard, Gentamycin	AI	IZ of standard, Streptomycin	AI
Lemon	<i>Malassezia furfur</i>	100%	50mm	16.5 mm	3.03	17 mm	2.94
Orange	<i>Malassezia furfur</i>	100%	20mm	16.5 mm	1.212	17 mm	1.17

Here, IZ= Inhibition zone (in mm) including the diameter of disc (6mm), AI (Activity index)

Table - 2: MIC of lemon oil against *Malassezia furfur*

Test strain	Different concentrations of lemon Oil used in $\mu\text{l ml}^{-1}$	Growth visually inspected in different concentrations of oil
<i>Malassezia furfur</i>	0.1	+4
	0.3	+3
	0.5	+2
	0.7	+1
	0.9	0
	1.1	0
	1.3	0
	1.5	0
	1.7	0
	1.9	0
Control without oil		100% growth

Growth was scored in the following manner:-

4+, growth comparable to that of the oil free control; 3+, growth approximately 75% that of the control; 2+, growth approximately 50% that of the control; 1+, growth 25% or less that of the control; and 0, no visible growth Here, μl represents the concentration of oil in per ml of brain heart infusion agar media

Table - 3: MIC of orange oil against *Malassezia furfur*

Test strain	Different concentrations of orange Oil used in $\mu\text{l ml}^{-1}$	Growth visually inspected in different concentrations of oil
<i>Malassezia furfur</i>	0.1	+4
	0.3	+4
	0.5	+4
	0.7	+4
	0.9	+3
	1.1	+3
	1.3	+3
	1.5	+3
	1.7	+2
	1.9	+2
	2.1	+1
	2.2	0
	2.4	0
	2.6	0
Control without oil		100% growth

Growth was scored in the following manner:-

4+, growth comparable to that of the oil free control; 3+, growth approximately 75% that of the control; 2+, growth approximately 50% that of the control; 1+, growth 25% or less that of the control; and 0, no visible grow

obtained against lemon and orange oil at 100% concentration of pure oil was 50 and 20 mm respectively by disc diffusion method. *M. furfur* was found to be resistant against clotrimazole and ketoconazole. Other reference antibiotics i.e gentamycin and streptomycin showed inhibition zone of 16.5 and 17 mm respectively. According to our study by comparing with the reference drugs, lemon and orange oil was found to be more effective in inhibiting the growth of *M. furfur*. Results of MIC of lemon and orange oils against *M. furfur* are summarized in Table 2,3. The results show that the lemon and orange oil exhibited inhibitory action at 0.8 and 2.2 $\mu\text{l ml}^{-1}$ concentrations against *M. furfur*. Both the oils can be used for the development of natural antifungal agent against *Pityriasis versicolor* infections.

Discussion

The present results suggest that lemon and orange oils exhibits strong antifungal activity. This is in agreement with the findings of (Valero and Frances, 2006) who reported narrow and wide ranges of antifungal activities. Essential oils derived from many plants are known to possess antifungal activities (Valero and Frances, 2006). Their mechanism of action appears to be predominantly on the fungal cell membrane, disrupting its structure causing leakage and cell death; blocking the membrane synthesis; inhibition of the spore germination, fungal proliferation and cellular respiration (Harris, 2002). Because of high volatility and lipophilicity of the essential oils, they are readily attached to penetrate into the cell membrane to exert their biological effect (Jayaprakasha *et al.*, 2005).

The results showed the efficacy of lemon oil on the animal skin infections caused by *Trichophyton rubrum* and *M. gypseum*. In our study, lemon oil exhibited the strong antimycotic activity against *M. furfur*. In screening of lemon oil, the diameter of inhibition zone by disc-diffusion method was found to be 50 mm. MIC of oil obtained by microdilution method was 0.8 $\mu\text{l ml}^{-1}$. Our work is in agreement with the observations of Silvia *et al.* (2008) in which screening of lemon grass oil against *Candida* species was studied and it was found as an effective antifungal agent. The diameter of inhibition zone against *Candida* species was found to be more than 40 mm. Our work also coincide with the previous findings of Ezzat (2001) which revealed that the essential oil of *Citrus lemon* at 200 μl concentration showed diameter of inhibition zone 37 mm against *Candida albicans* 10261. According to Rusenova and Parvano (2009) *Citrus lemon* does not show antimicrobial activity by disc-diffusion method against both yeasts, *Candida albicans* and *M. pachydermatis*. These results are not in agreement with our studies which showed that antifungal activity by disc-diffusion and minimum inhibitory concentration by microdilution method, lemon oil was found to be effective in inhibiting the growth of *M. furfur* at a low concentrations of oil. Lemon oil, is also preferentially more active against Gram-positive bacteria as compared to Gram-negative bacteria. The antimicrobial activity of lemon oil is due to the presence of terpenoids- α -pinene, camphene, β -pinene, sabinene, myrcene, terpinene, β -bisabolene, limonene, trans- α -bergamotene, nerol and neral (Ahmed and Beg, 2001).

Essential oils from citrus offer the potential for all natural antimicrobials for use in improving the safety of organic or all natural foods (Bryan *et al.*, 2008). Shahi *et al.* (2003) found that the MIC of oil of *Citrus sinensis* were in the range of 0.7-1 $\mu\text{l ml}^{-1}$ against *Candida albicans*, *A. flavus*, *A. fumigatus*, *A. niger*, *A. ustus*, *Epidermophyton floccosum*, *Microporum audouinii*, *M. canis*, *M. gypseum*, *M. nanum*, *Rhizopus nigricans*, *Trichophyton tonsurans*, *T. rubrum*, *T. mentagrophytes* and *T. violaceum*. The present investigation suggests that the essential oils exhibit narrow or wide range of antifungal activity, which may prove useful in the development of effective antifungal substances. Souza *et al.* (2005) reported MIC of lemon oil at 0.5% concentration against mould strains i.e *Fusarium* spp, *Penicillium* spp, *Aspergillus niger*, *A. flavus* and *Rhizopus* spp.

In our findings, MIC of lemon oil against *M. furfur* was $0.8 \mu\text{l ml}^{-1}$ i.e very much close to that concentration. The small differences may be due to local environmental and climatic conditions and variation in choice of microorganism tested.

Our results of orange (*Citrus sinensis*) and lemon (*Citrus lemon*) are also similar to the work of Viuda-Martos *et al.* (2008) which suggests that both the oils were effective in inhibiting the growth of *A. niger*, *A. flavus*, *Penicillium chrysogenum* and *P. verrucosum*. The hydrodistilled oil of lemon and orange was also found to be more sensitive against *Penicillium digitatum* (Caccioni *et al.*, 1998). Antimicrobial activity of Rutaceae family species have been observed on dermatophytes (Lima, 1996) and on some moulds (Megalla *et al.*, 1980). Essential oils of some plants have recently been proven to be successful eco-friendly, bio-control agent (Chutia *et al.*, 2006). Many authors have reported antimicrobial, antifungal, antioxidant and radical-scavenging properties of essential oils (Sawamura *et al.*, 2005; Sokovic and Griensven, 2006).

The present study clearly suggests that the oils extracted from the waste product (peels) of *C. sinensis* and *C. lemon* hold good promise as an antifungal agent, which could be used in therapeutic remedy against human pathogenic fungi on account of its various antifungal properties, viz. strong fungicidal action, long-shelf life, withstand heavy inoculum density, thermostable, broad range of antifungal activity and absence of any adverse effects. The waste product of *C. sinensis* and *C. lemon* can be used for the development of a potential source of effective and economically viable herbal antifungal against pityriasis versicolor (fungal skin infection) after undergoing successful clinical trials.

Acknowledgements

This work was financially supported by Department of Science and Technology (Government of India) Delhi and we thank Dr Rakesh Meena, E.S.I.C Hospital, Jaipur for providing constant support and guidance during the collections of clinical samples.

References

- Ahmed, I. and Beg, A.Z.: Antimicrobial and phytochemical studies on Indian medicinal plants against multidrug resistant human pathogens. *J. Ethnopharmacol.*, **74**: 113-123 (2001).
- Alex, L.A., Korch, C., Selitrennikoff, C.P. and Simon, M.I.: COS1, a two-component histidine kinase that is involved in hyphal development in the opportunistic pathogen *Candida albicans*. *Proc. Natl. Acad. Sci.*, **95**: 7069-7073 (1998).
- Bauer, A.W., Kirby, W.M., Sherris, J.C. and Turck, M.: Antibiotic susceptibility testing by standardized single disc method. *Am. J. Clin. Pathol.*, **44**: 493-496 (1996).
- Bektas, T., Dimitra, D., Munewer, S., Moschos, P. and Atalay, S.: In vitro antimicrobial and antioxidant activities of the essential oils and various extracts of *Thymus eigi* M. Zohary et P.H. Davis. *J. Agric. Food Chem.*, **52**: 1132-1137 (2004).
- Boussida, S., Boudaya, S., Ghorbel, R., Meziou, T.J., Markkekehi, S. and Turki, H.: *Pityriasis versicolor* in children: a retrospective study of 164 cases. *Ann. Dermatol. Venereol.*, **125**: 581-584 (1998).
- Bryan, C.A.O., Crandall, P.G., Chalova, V.I. and Ricke, S.C.: Orange essential oils antimicrobial activities against *Salmonella* spp. *J. Food Sci.*, **63**: 264-267 (2008).
- Burt, S.: Essential oils: their antibacterial properties and potential applications in foods-A review. *Int. J. Food Microbiol.*, **94**: 223-53 (2004).
- Caccioni, D.R.L., Guizzardi, M., Biondi, D.M., Agatino, R. and Ruberto, G.: Relationship between volatile oils of components of citrus fruits, essential oils and antimicrobial action on *Penicillium digitatum* and *Penicillium italicum*. *Int. J. Food Microbiol.*, **43**: 73-79 (1998).
- Davies, J.: In activation of antibiotic and the dissemination of resistance genes. *Sci*; **264**: 375-382 (1994).
- Ezzat, S.M.: In vitro inhibition of *Candida albicans* growth by plant extracts and essential oils. *World J. Microbiol. Biotechnol.*, **17**: 757-759 (2001).
- Faergemann, J. and Freidriksson, T.: Experimental infections in rabbits and human with *Pityriasis orbicularis* or *Pityriasis ovale*. *J. Invest Dermatol.*, **77**: 314-318 (1981).
- Harris, R.: Progress with superficial mycoses using essential oils. *Int. J. Aromatherapy*, **12**: 83-91 (2002).
- Hay, R.J. and Moore, M.: Book of Dermatology. 6th Edition. In: R.H. Champion, J.L. Burton, D.A. Burns (Eds), Blackwell Science Oxford. England. pp. 1277-1290 (1998).
- Jayaprakasha, G.K., Jaganmohan Rao, L. and Sakariah, K.K.: Chemistry and biological activities of *C. longa*. *Trend. Food Sci. Technol.*, **16**: 533-548 (2005).
- Kordali, S., Kotan, R., Mavi, A., Cakir, A., Ala, A. and Yildirim, A.: Determination of the chemical composition and antioxidant activity of essential oil of *Artemisia dracuncululus* and antifungal and antibacterial activities of Turkish *Artemisia absinthium*, *Artemisia dracuncululus*, *Artemisia santonicum*, and *Artemisia spicigera* essential oils. *J. Agric. Food Chem.*, **53**: 9452-9458 (2005).
- Lima, E.O.: Estudo das dermatofitoses em João Pessoa-PB e da atividade de plantas medicinais contra alguns dos fungos isolados. PhD Thesis, University of Sao Paulo; Sao Paulo (1996).
- Marcon, M.J. and Powell, D.A.: Human infection due to *Malassezia* spp. *Clin. Microbiol. Rev.*, **5**: 101-119 (1992).
- Megalla, S.E., Ross, S.A. and Kektani, N.E.: Antimicrobial activity of some Egyptian aromatic plants. *Fitoterapia*, **51**: 201-206 (1980).
- Paranagama, P.A., Abeysekera, K.H.T., Abeywickrama, K. and Nugaliyadde L.: Fungicidal and anti-aflatoxinigenic effects of the essential oil of *Cymbopogon citratus* (DC.) Stapf (lemongrass) against *Aspergillus flavus* Link. Isolated from stores rice. *Lett. Appl. Microbiol.*, **37**: 86-90 (2003).
- Provine, H. and Hadley, S.: Preliminary evaluation of a semisolid agar antifungal susceptibility test for yeasts and molds. *J. Clin. Microbiol.*, **38**: 537-541 (2000).
- Rios, J.L., Recio, M.C. and Vilar A.: Screening methods for natural products with antimicrobial activity: a review of the literature. *J. Ethnopharmacol.*, **23**: 127-49 (1988).
- Rippon, J.W.: Dermatophytosis and Dermatophytomycosis. In: Medical Mycology. 2nd Edition. Publisher. W.B. Saunders, Philadelphia. pp. 154-159 (1982).
- Rusenova, N. and Parvano, P.: Antimicrobial activities of twelve essential oils against microorganisms of veterinary importance. *Trakia J. Sci.*, **7**: 37-43 (2009).
- Sawamura, M., Tu, N.M., Yu, X. and Xu, B.: Volatile constituents of the peel oils of several sweet oranges in China. *J. Essential Oil Res.*, **17**: 2-6 (2005).
- Shahi, S.K., Patra, M. and Dikshit, A.: Utilization of pericarp of *Citrus sinensis* oil for the development of natural antifungal against nail infection. *Curr. Sci.*, **84**: 1512-1514 (2003).
- Schmidt, A.: *Malassezia furfur*: A fungus belonging to the physiological skin flora and its relevance in skin disorders. *Cuts.*, **59**: 21-24 (1997).
- Singh, N.: Trends in the epidemiology of opportunistic fungal infections: predisposing factors and the impact of antimicrobial use practices. *Clin. Infect. Diseases*, **33**: 1692-1696 (2001).
- Sofowora, A.: Introduction to medicinal plants and traditional medicine. *Spectrum Books limited*, **2**: 8-76 (1993).
- Silva-Lizama E.: Tinea versicolor. *Int. J. Dermatol.*, **34**: 611-617 (1995).
- Silva, C.de.Bda, Guterres, S.S. and Weisheimer, V.: Antifungal activity of the lemon grass oil and citral against *Candida* spp. *Braz. J. Infect. Diseases*, **12**: 63-66 (2008).
- Sokovic, M. and Griensven Leo, J.L.D Van.: Antimicrobial activity of essential oil and their components against the three major pathogens of cultivated button mushroom *Agaricus bisporus*. *Eur. J. Plant Pathol.*, **116**: 211-224 (2006).
- Valero, M. and Francés, E.: Synergistic bactericidal effect of carvacrol, cinnamaldehyde or thymol and refrigeration to inhibit *Bacillus cereus* in carrot Broth. *Food Microbiol.*, **23**: 68-73 (2006).
- Viuda-Martos, M; Ruiz-Navajas, Y; Fernandes-Lopez, J. and Perez-Alvarez, J. Antifungal activity of lemon (*Citrus lemon* L.), mandarin (*Citrus reticulata* L.), grapefruit (*Citrus paradisi* L.) and orange (*Citrus sinensis* L.) essential oils. *Food Control*, **19**: 1130-1138 (2008).